



Role of the Nervous System in Tumor Angiogenesis

Nyanbol Kuol¹ · Lily Stojanovska¹ · Vasso Apostolopoulos¹ · Kulmira Nurgali^{1,2} 

Received: 18 November 2017 / Accepted: 19 February 2018 / Published online: 4 March 2018
© Springer Science+Business Media B.V., part of Springer Nature 2018

Abstract

The development of cancer involves an intricate process, wherein many identified and unidentified factors play a role. Tumor angiogenesis, growth of new blood vessels, is one of the major prerequisites for tumor growth as tumor cells rely on adequate oxygen and nutrient supply as well as the removal of waste products. Growth factors including VEGF orchestrate the development of angiogenesis. In addition, nervous system via the release of neurotransmitters contributes to tumor angiogenesis. The nervous system governs functional activities of many organs, and, as tumors are not independent organs within an organism, this system is integrally involved in tumor growth and progression via regulating tumor angiogenesis. Various neurotransmitters have been reported to play an important role in tumor angiogenesis.

Keywords Nervous system · Neurotransmitters · Neuropeptides · Neuro-cancer interaction · Angiogenesis · Cancer

Introduction

New growth in the vascular network (angiogenesis) is a normal physiological phenomenon that tumors utilize to aid in their growth, proliferation and metastatic spread. Angiogenesis involves migration and division of endothelial cells, generation of new basement membrane, arrangement into tubular structures and coverage by pericytes. Angiogenesis is regulated by a plethora of pro- and anti-angiogenic molecules such as, interleukin (IL)-8, tumor necrosis factor (TNF)- α , vascular endothelial growth factor (VEGF), transforming growth factor (TGF)- α , TGF- β , angiogenin, platelet-derived growth factor (PDGF) and fibroblast growth factor (FGF) [1, 2]. The level of angiogenic factors in tissues reflects the aggressiveness of tumor cells which play a significant role in prognostic outcomes [3, 4]. In cancer, the balance between pro- and anti-angiogenic factors is lost, resulting in uncontrolled angiogenesis with irregular blood vessels lacking a clear hierarchal arrangement [1, 5]. As a

consequence, anti-angiogenic therapies (in particular anti-VEGF) have been approved for cancer treatment [4, 6–8]. The interaction between VEGF with its receptor, VEGFR2, is responsible for the majority of the angiogenic stimulatory signals in vivo, however, their therapeutic value for long-term patient survival is relatively modest [3].

In addition to these factors, the impact of the tumor micro-environment in tumor angiogenesis has attracted much interest in recent years as another regulator of angiogenesis [9–12]. Furthermore, the role of the nervous system has also surfaced as one of the major contributors to cancer progression through the regulation of tumor angiogenesis via release of neurotransmitters. The nervous system governs functional activities of many organs, and, as tumors are not independent organs within an organism, this system is integrally involved in tumor growth and progression [13, 14]. Here we present an overview of the nervous system role in tumor angiogenesis.

Neurotransmitters Influencing Tumor Angiogenesis

Neurotransmitters are group of neurological chemical messengers synthesized by neurons and secreted at nerve terminals where they transmit signals to target cells through binding to their receptors. Studies have demonstrated that various cancers express receptors for different neurotransmitters which have been identified to play essential role in the control of tumor angiogenesis (Table 1, Fig. 1).

Vasso Apostolopoulos and Kulmira Nurgali contributed equally to this work.

✉ Kulmira Nurgali
Kulmira.Nurgali@vu.edu.au

¹ Centre for Chronic Disease, College of Health and Biomedicine, Victoria University, Melbourne, Australia

² Department of Medicine Western Health, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Regenerative Medicine and Stem Cells Program, Australian Institute for Musculoskeletal Science (AIMSS), Melbourne, Australia

Table 1 Neurotransmitters influencing tumor angiogenesis

Neurotransmitters		Receptor	Type of cancer	Model	Mechanism/pathway	Ref.
NE	Breast cancer	β 2-AR		MCF-7, MDA-453, and MDA-231 cell lines, subcutaneous injection of 4 T1 cells in BALB/c mice	β 2-AR expression is elevated in MDA-453, decreased in MCF-7 and intermediate in MDA-231 cells. Administration of β -AR agonist, isoproterenol upregulates Jagged 1 expression and enhances tumor microvessel density via NE-induced β 2-AR/PKA/mTOR pathway in vivo.	[15]
				HT-29 and CT26 cells in vitro and subcutaneous injection of HT-29 cells in nude mice and CT26 cells in BALB/c mice	Activation of β 2-AR by NE enhances expression of VEGF, IL-8 and IL-6 in vitro and in vivo \rightarrow stimulation of tumor angiogenesis via β -AR cAMP-PKA signaling pathway.	[16]
	Melanoma			B16-F1 cells in vitro and subcutaneous injection in the flanks of C57BL/6 mice		[17]
	Lung adenocarcinoma			A549 cells in vitro		
DA	Ovarian cancer	DR1 & DR2		SKOV3p 1, HeyA8 cells in vitro and intraperitoneal injection of these cells in a chronic stress C57BL/6 mouse model	Activation of DR2 mediates inhibitory effect of DA on tumor angiogenesis cAMP-PKA signaling pathway.	[18, 19]
	Gastric cancer			Human gastric cancer tissues, subcutaneous injection of Hs746T cells in nude mice, MNNG-induced gastric cancer in rats	DA suppresses gastric cancer growth by inhibition of VEGF-stimulated angiogenesis. In both human gastric cancer and MNNG-induced animal model DA is depleted. Suppression of VEGFR-2 phosphorylation in endothelial cell \rightarrow inhibition of angiogenesis.	[20]
	Lung cancer			Orthotopic injection of LLC1 cells in C57BL/6 mice and A549 cells in SCID mice	Administration of DR2 agonists inhibits in vivo lung tumor progression via suppressing angiogenesis and reducing myeloid-derived suppressor cells infiltration.	[21]
GABA	Cholangiocarcinoma	GABA		H-69, Mz-ChA-1, HuH28, and TFK-1 cells, subcutaneous injection of Mz-ChA-1 cells in BALB/c mice	GABA _A , GABA _B , and GABA _C receptors were expressed by cells in vitro which inhibit cell growth and proliferation via IP3/cAMP, PKA phosphorylation, and ERK1/2 dephosphorylation. GABA \downarrow tumor size and VEGF-A/C expression in vivo.	[22]
5-HT	Ovarian cancer	GABA _A		OVCAR-3 cells in vitro	\uparrow Level of GABARBP inhibited VEGF expression and \downarrow HIF-1 α protein via PI3K-mTOR-4E-BP1 signaling pathway in vitro.	[23]
	Colon cancer	5-HT receptor		Subcutaneous injection of MC-38 cells in <i>Tph1</i> ^{-/-} mice	5-HT regulates angiogenesis by reducing MMP12 expression in TAMs, thus affecting the production of circulating angiostatin.	[24]
Glu	Breast cancer	mGluR1 on endothelial cells		4 T1 cells injected into the mammary fat pads of BALB/c mice	\downarrow mGluR1 expression results in \downarrow angiogenesis in vivo.	[25]
	Melanoma	GRM1		UACC903-G2, UACC903-G4, C8161-G21, C81-61-G6, and C81-61-G7 cells, subcutaneous injection of these cells into each flank of nude mice	In vitro \uparrow expression of GRM1 \rightarrow \uparrow expression of IL-8 and VEGF via the AKT-mTOR-HIF1 signaling pathway activation. In vivo \uparrow expression of GRM1 \rightarrow larger melanoma tumors.	[26]
ACh	Lung cancer	α 7-nAChRs		Human NSCLC A549 and H157 cell lines	Nicotine increases HIF-1 & VEGF expression. Nicotine mediates tumor angiogenesis through PI3K/Akt and ERK1/2 signalling pathway.	[27]
	Colon cancer			Subcutaneous injection of HT-29 cells in BALB/c mice	Administration of nicotine \uparrow VEGF expression \rightarrow \uparrow microvessel densities and angiogenesis via stimulation of β 2-AR.	[28]
	Breast cancer	mAChR				[29]

Table 1 (continued)

Neurotransmitters	Receptor	Type of cancer	Model	Mechanism/pathway	Ref.
NPY	Y5R	Mammary adenocarcinoma	Intradermal injection of MCF-7 and MCF-10A cells in nude mice, intradermal injection of LMM3 cells in BALB/c mice	mAChR activation promotes VEGF-A production and neovascularisation in breast cancer models.	[30]
		Breast cancer	Subcutaneous injection of LMM3 cells in BALB/c mice 4 T1 cell line	↑ Expression of VEGF by activation of M1 and M2 mAChRs via arginine metabolic pathway. Activation of NPY ↑ the expression and secretion of VEGF → angiogenesis.	
NO	Y2R	Melanoma	Subcutaneous injection of B16F10 cells into C57BL/6 mice	Blockade of the Y2R inhibited tumor growth by ↓ tumor angiogenesis.	[33]
		Neuroblastomas	Human tissue	Y2R expression is observed in both tumor and endothelial cells.	[34]
		Breast cancer	MDAMB-231 cell and human invasive breast cancer tissues	NO induces the expression of VEGF-C in both breast cancer cell line and human tissues.	[35]
		Ovarian cancer	Cystic fluid samples and human tissues	The expression of iNOS correlates with the degree of tumor differentiation; level of intracyclic NO metabolite correlates with tumor stage.	[36]
		Gastric cancer	Human tissues (all stages)	NOS III protein is ↑ in both primary gastric tumors and lymph node metastases.	[37]

α7nAChR, α7 nicotinic acetylcholine receptor; ACh, acetylcholine; β2-AR, β2-adrenergic receptor; cAMP, cyclic adenosine monophosphate; AKT, serine/threonine kinase or protein kinase B; DA, dopamine; DR1 & DR2, dopamine receptor 1 & 2; ERK1/2, extracellular signal-regulated kinase; GABA, gamma-aminobutyric acid; GABARBP, GABA_A receptor-binding protein; GABA_{A,B&C}, gamma-aminobutyric acid receptor _{A,B&C}; Glu, glutamate; GRM1, glutamate receptor metabotropic 1; HIF-1α, hypoxia inducible factor-1α; 5-HT, 5-hydroxytryptamine (serotonin); iNOS, inducible nitric oxide synthase; IL-6, interleukin 6; IL-8, interleukin 8; mGluR1, metabotropic glutamate receptor 1; mAChRs, muscarinic acetylcholine receptors; M1 & M2, muscarinic 1 & 2 receptors; MMP12, matrix metalloproteinase 12; MNNG, N-methyl N'-nitro-N-nitrosoguanidine; mTOR, mammalian/mechanistic target of rapamycin; NE, norepinephrine; NO, nitric oxide; NOS, nitric oxide synthase; NPY, neuropeptide Y; NSCLC, non-small cell lung carcinoma; PI3K, phosphoinositide 3-kinase; 4E-BP1, phosphorylated 4E binding protein 1; PKA, protein kinase A; TAMs, tumor-infiltrating macrophages; VEGF, vascular endothelial growth factor; VEGFR, vascular endothelial growth factor receptor; Y2R & Y5R, neuropeptide Y receptor 2 & 5




Fig. 1 Neurotransmitter signalling pathways in cancer angiogenesis. Neuro-cancer communication is through the release of neurotransmitters activating different signalling kinases which promote cancer progression via angiogenesis. ACh, acetylcholine; β₂-AR, β₂-adrenergic receptor; cAMP, cyclic adenosine monophosphate; AKT, serine/threonine kinase or protein kinase B; DA, dopamine; DR, dopamine receptor; ERK_{1/2}, extracellular signal-regulated kinase; GABA, gamma-aminobutyric acid; GABA_{A&B}, gamma-aminobutyric acid receptor_{A&B}; Glu, glutamate; GRM1, glutamate receptor metabotropic 1; HIF-1, hypoxia inducible

factor 1; 5-HT, 5-hydroxytryptamine (serotonin); 5-HTR, 5-hydroxytryptamine receptor (serotonin); MMP12, matrix metalloproteinase 12; mTOR, mammalian/mechanistic target of rapamycin; nAChR, nicotinic acetylcholine receptor; NE, norepinephrine; NPY, neuropeptide Y; PI3, phosphoinositide 3; PI3K, phosphoinositide 3-kinase; 4E-BP1, phosphorylated 4E binding protein 1; PKA, protein kinase A; p70S6K, serine/threonine kinase; VEGF, vascular endothelial growth factor; Y5R, neuropeptide receptor

Catecholamines are a group of neurotransmitters that are synthesized from amino acid tyrosine. These neurotransmitters are intricately involved in the normal physiological response of fight or flight response during stress [38, 39]. Epinephrine and norepinephrine released during chronic stress play an important role in tumorigenesis via regulation of angiogenesis through β-adrenergic signaling. The β-adrenergic signaling pathway is involved in regulation of cancer initiating factors such as apoptosis, DNA damage repair, inflammation, cellular immune response, angiogenesis and epithelial-mesenchymal transition. Numerous in vitro and animal studies have demonstrated that epinephrine and

norepinephrine acting on their receptors expressed on tumor cells, stimulate angiogenesis via increased VEGF synthesis [16, 38–41] through the cAMP-PKA signaling pathway [40]. In fact, activation of the β-adrenergic signaling pathway in primary mammary tumors has been shown to elevate tumor-associated macrophages (TAMs) expressing *veg*f gene which enhances angiogenesis [42]. Moreover, in some breast cancer cell lines, direct activation of β-adrenergic signaling can amplify expression of VEGF and cytokines, IL-6, and IL-8 that stimulate tumor angiogenesis [43]. Jagged 1 is essential factor mediating Notch signaling which regulates tumor angiogenesis through β₂-AR-PKA-mTOR pathway.

Upregulation of Jagged 1 in breast cancer patients correlates with poor prognosis [44, 45]. Knockdown of Jagged 1 by siRNA in MDA-231 breast cancer cells inhibits Notch signaling in endothelial cells and impairs tumor angiogenesis induced by norepinephrine [15].

In contrary, dopamine inhibits angiogenesis by down-regulation of VEGFR-2-mediated signaling pathway in both tumor endothelial and endothelial progenitor cells through D₂ dopamine receptors (DR2) [38, 39, 46, 47]. Furthermore, in mouse models of breast cancer induced by MCF-7 cell line and colon cancer induced by HT29 cell line, dopamine administration in combination with anti-cancer drugs (eg. doxorubicin and 5-fluorouracil) impairs tumor growth and improves survival outcome [48]. However, dopamine effect was found to have no direct impact on tumor growth and survival but by inhibiting tumor endothelial cell proliferation and migration via the suppression of VEGFR-2 and mitogen-activated protein kinase as demonstrated in vitro [48]. In tissues from gastric cancer patients and in rats with chemically-induced as well as mice with Hs746T cell-induced gastric cancer, administration of dopamine decelerates tumor growth by suppressing angiogenesis via inhibition of VEGFR-2 phosphorylation in endothelial cells [20]. This concurs with results obtained in ovarian cancer mouse models induced by systemic injection of SKOV3ip1 and HeyA8 cells in which exogenous administration of dopamine inhibits angiogenesis by a stimulation of DR2, however stimulation of DR1 stabilizes tumor blood vessels via cAMP-PKA signaling pathway [18].

Acetylcholine and Nicotine Nicotinic acetylcholine receptors (nAChRs) can have either stimulatory or inhibitory effect on the production and release of angiogenic factors [49]. Indeed, the expression of VEGF, TGF- β , FGF and PDGF in endothelial cells is increased by nicotine [50–53]. Nicotine-mediated angiogenesis via activation of $\alpha 7$ and $\alpha 9$ -nAChRs is cell-type specific, e.g. in lung cancer cells angiogenesis is promoted via activation of $\alpha 7$ -nAChRs [53, 54], whereas in breast tumors overexpression of $\alpha 9$ -nAChRs [55] stimulates release of pro-angiogenic factors [56]. In colon tumor tissues from HT-29 cell-bearing BALB/c mice, VEGF expression is elevated by nicotine which correlates with enhanced microvessel density [28]. The molecular pathways of nicotine-induced angiogenesis have been extensively reviewed [57]. The role of muscarinic acetylcholine receptors (mAChRs) in tumor angiogenesis is not well understood, however administration of autoantibodies against mAChRs in mouse models of breast cancer (Table 1) mediates tumor angiogenesis via activation of mAChRs through release of VEGF-A [29]. In addition, in BALB/c mice bearing LMM3 mammary adenocarcinoma cells, administration of muscarinic agonist, carbachol, in the presence or absence of various muscarinic antagonists shows an increase in VEGF expression [30, 58].

Furthermore, tumor macrophages stimulate angiogenesis via activation of M1 and M2 mAChRs which trigger arginine metabolic pathway [30].

Y-Aminobutyric Acid (GABA), Neuropeptide Y (NPY), Nitric Oxide (NO) and Serotonin have varying effects on angiogenesis and tumor progression. In a mouse model of cholangiocarcinoma, GABA inhibits VEGF-A/C, decreases cell proliferation and tumor mass [22]. NPY enhances the expression of VEGF and its secretion promoting angiogenesis and breast cancer progression [31]. The suggested mechanism by which NPY induces angiogenesis is by its influence on endothelial cells dependent on endothelial nitric oxide synthase (eNOS) activation and partly on VEGF signaling pathway. The release of NO results in endothelial activation inducing tumor cells lysis [59], although NO can also promote tumor growth and metastasis by enhancing angiogenesis [36, 59–65]. For instance, NO increases VEGF-C and nitrite/nitrate production in MDA-MB-231 breast cancer cells and high levels of nitrotyrosine correlate with increased VEGF-C, lymph node metastasis, reduced disease-free and overall survival in invasive breast carcinoma [35]. The expression of iNOS and VEGF in colorectal cancer correlates with enhanced intratumor micro-vessel density suggesting that NO can promote tumor angiogenesis [60]. In gastric cancer, overexpression of NOS III via abnormal activation of sequence-specific DNA-binding protein (Sp1) correlates with enhanced micro-vessel density and poor survival [37]. Serotonin has also been implicated in tumor angiogenesis. In C57BL/6 mice bearing MC-38-induced tumors, serotonin regulates angiogenesis by plummeting matrix metalloproteinase 12 (MMP12) expression (eg. [66]) in macrophages infiltrating the tumor, as well as reducing angiostatin (an endogenous inhibitor of angiogenesis) levels [24].

Glutamate is an excitatory neurotransmitter that regulates synaptic and cellular activity via binding to its receptors including metabotropic glutamate receptors (mGluRs). The expression of mGluRs has been implicated in tumor angiogenesis as noted in mouse models of melanoma and breast cancer [25, 26, 67]. As such, decreased activity of mGluR1 inhibits angiogenesis in an orthotopic breast cancer (4 T1) model suggesting that mGluR1 acts as a pro-angiogenic and pro-tumorigenic factor [25]. Likewise, in an experimental non-small cell lung cancer in A549-bearing nude mice, inhibition of mGlu1 receptor with BAY36–7620 led to suppression of angiogenesis via inhibiting AKT/HIF-1 α /VEGF signaling pathway [68]. Similarly, high expression of glutamate receptor GRM1 in several human melanoma cell lines (Table 1) leads to increased expression of IL-8 and VEGF via activation of the AKT/mTOR/HIF1 signaling pathway [26].

Table 2 Other factors influencing tumor angiogenesis

Factors	Type of cancer	Model	Mechanism/pathway	Ref.
ANG	Breast cancer	Human tissues	The level of ANG correlates with clinical progression. ANG derived from tumors activates angiogenesis via suppression of miR-543-2p.	[69]
	Bladder cancer	Human tissues, T24, UROtsa and HeLa cells subcutaneously injected in athymic BALB/c (nu/nu) mice	↑ ANG expression correlates with high grade, and muscle-invasive tumors via ERK 1/2 and MMP2. Downregulation of ANG inhibits tumor angiogenesis via AKT/GSK3 β / mTOR pathways.	[70, 71]
TNF- α	Lung cancer	LLC1 cells subcutaneously injected in wild type, p75 knockout (KO) and double p55KO/p75KO mouse xenograft models	Tumor growth ↓ in both LLC and B16 p75KO mice. Decreased tumor growth correlates with ↓ VEGF expression and capillary density via TNFR2/p75.	[72]
	Melanoma	B16 cell subcutaneously injected in C57BL/6 mice. Wild type, p75 knockout (KO) and double p55KO/p75KO mouse tumor xenograft models		
TGF- β	Colon cancer	Human tissues, FET α /DNRII cell	TGF- β signaling is inversely correlates with the expression of VEGF-A in tissues. TGF- β ↓ VEGF-A expression via ubiquitination and deterioration in a PKA- and Smad3-dependent and Smad2-independent pathways in vitro.	[73]
BDNF	Chondrosarcoma	JJ012 cell line, JJ012 cells subcutaneously injected in CB17-SCID mice	The expression of BDNF and VEGF correlates with tumor grade. BDNF knockdown ↓ angiogenesis and tumor growth in mouse model. BDNF ↑ expression of VEGF and stimulates angiogenesis via the TrkB receptor, PKC α , PLC γ and HIF-1 α signaling pathways.	[74]
FGF	Mammary cancer	Mouse 66c14 mammary carcinoma and inguinal mammary fat pad injection in BALB/c mice	In tumor cells suppression of FGFR signaling inhibits expression of VEGF-C and induces VEGFR-3, netrin-1, prox1 and integrin α 9 expression.	[75]
	Glioma	Rat C6 glioma cancer cells injected subcutaneously into rats		
EGFR	HNSCC	Human tissues, CAL27 cells subcutaneously injected in nude mice	In human tissues, ↑ EGFR correlates with ↑ HIF-1 α and microvessel density. EGFR inhibitors ↓ the regulation of HIF-1 α & Notch1 → ↓ angiogenesis and tumor size.	[76]
NGF	Breast cancer	MDA-MB-231 cells subcutaneously injected into SCID mice	NGF ↑ the release of VEGF in breast cancer cells and mediates angiogenic effect via the activation of PI3K-Akt, ERK, MMP2 and NO synthase pathways.	[77]
HGF	ESCC	Serum samples, human tissues, HKESC-1, HKESC-2 and SLMT cells	In tissues, ↑ level of HGF correlates with tumor metastasis and poorer survival. In serum samples, ↑ HGF level correlated with expression of VEGF and IL-8. HGF stimulates cells to express VEGF and IL-8 in vitro via extracellular signal-regulated kinase signaling pathways.	[78]
	Prostate cancer	Castration-resistant prostate cancer blood samples and PC3 cell line	HGF levels ↑ in both blood samples and cell line.	[79]

AKT, serine/threonine kinase or protein kinase B; ANG, angiogenin; BDNF, brain-derived neurotrophic factor; EGFR, epidermal growth factor receptor; ESCC, esophageal squamous cell carcinoma; ERK_{1/2}, extracellular signal-regulated kinase; FGF, fibroblast growth factor; FGFR, fibroblast growth factor receptor; GSK3 β , glycogen synthase kinase 3 β ; HNSCC, head and neck squamous cell carcinoma; HGF, hepatocyte growth factor; HIF-1 α , hypoxia inducible factor 1 α ; IL-8, interleukin-8; MMP2, matrix metalloprotease 2; mTOR, mammalian/mechanistic target of rapamycin; NGF, nerve growth factor; NO, nitric oxide; PI3K, phosphoinositide 3-kinase; PKA, protein kinase A; PKC α , protein kinase C alpha; PLC γ , phospholipase C γ ; TGF- β , transforming growth factor beta; TNF- α , tumor necrosis factor alpha; TNFR2/p75, tumor necrosis factor receptor 2/neurotrophin receptor; TrkB, tropomyosin related kinase B; VEGF, vascular endothelial growth factor

Hence, these studies clearly demonstrate involvement of neurotransmitters in tumor angiogenesis; however, most of the studies have been performed mainly in animal models

and cell lines. Understanding their relevance to human pathology may aid in the development of better anti-angiogenic therapies.

Other Factors Influencing Tumor Angiogenesis

Nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), angiogenin (ANG), FGF, TNF- α , TGF- β , hepatocyte growth factor (HGF) and epidermal growth factor receptor (EGF) are important signaling molecules promoting angiogenesis (Table 2, Fig. 2). NGF is a neurotrophic factor that is upregulated in tumor microenvironment of various cancers including breast cancer [77]. NGF, secreted by MDA-MB-231 breast cancer cells, stimulates angiogenesis in vivo after

injection of these cells subcutaneously to immunodeficient mice and enhances endothelial cell proliferation, invasion, migration and tubule formation in vitro [77]. Furthermore, NGF enhances secretion of VEGF by breast cancer cells; in vivo administration of anti-VEGF antibody inhibits its angiogenic capacity [77]. In human glioma microvascular endothelial cells, NGF mediates tumor angiogenesis by interaction with $\alpha 9 \beta 1$ integrin [80–83]. Another neurotrophic factor, BDNF has been shown to play a role in tumor angiogenesis. For instance, in chondrosarcoma patients, BDNF and VEGF protein expression is significantly higher which is correlated with




Fig. 2 Growth factors intracellular signalling pathways in cancer angiogenesis. The binding of growth factors to their respective receptors (eg. EGF to EGFR) activates multiple kinase pathways which are involved in cancer angiogenesis. AKT, serine/threonine kinase or protein kinase B; ANG, angiogenin; BDNF, brain-derived neurotrophic factor; CEBPB, CCAAT/enhancer-binding protein beta; EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; ERK_{1/2}, extracellular signal-regulated kinase; FGF, fibroblast growth factor; FGFR, fibroblast growth factor receptor; GSK3 β , glycogen synthase kinase 3 beta; HGF, hepatocyte growth factor; c-Met, hepatocyte growth factor receptor; HIF-1 α , hypoxia inducible factor 1 alpha; ICAM-1, intercellular

adhesion molecule-1; MAPK, mitogen activated protein kinase; MEK_{1/2}, MAPK/ERK kinase; MMP2, matrix metalloproteinase 2; mTOR, mammalian/mechanistic target of rapamycin; NGF, nerve growth factor; NF- κ B, nuclear factor-kappa B; NOS, nitric oxide synthase; PI3K, phosphoinositide 3-kinase; PKC- α , protein kinase C alpha; PLC- γ , phospholipase C-gamma; POU2F1, POU domain class 2 transcription factor 1; RAF, mitogen activated protein kinase; RAS, mitogen activated protein kinase; Tie2, angiopoietin receptor 2; TrkA, tropomyosin related kinase A; TrkB, tropomyosin related kinase B; VEGF, vascular endothelial growth factor

tumor stage [74]. Furthermore, BDNF knockdown decreases the expression of VEGF and abolishes angiogenesis in in vitro studies and animal models of chondrosarcoma [74].

In addition to neurotrophic factors, angiogenic factor ANG is upregulated in number of cancers [84–86] and is associated with worse clinical prognosis in urothelial carcinoma patients [87]. ANG regulates tumor angiogenesis via activation of endothelial and smooth muscle cells triggering various molecular pathways involved in the initiation of angiogenesis (Fig. 2) [69–71, 88]. Elevated expression of ANG associates with high grade and muscle-invasive human bladder tumors involving increase p-ERK1/2 and MMP2 expression [70]. Similarly, downregulation of ANG inhibits tumor angiogenesis via AKT/GSK3 β /mTOR pathways [71]. FGF is involved in angiogenesis by suppressing VEGF-C expression and stimulating expression of pro-lymphangiogenic factors including integrin α 9, VEGFR-3, prox1 and netrin-1 [75]. In fact, blocking of FGF2 with anti-FGF2 monoclonal antibody results in impaired angiogenesis of B16-F10 cell induced melanoma in mice [89]. In addition, TNF- α binding to TNFR1/p55 and TNFR2/p57 receptors has been implicated in the secretion of cytokines and pro-angiogenic factors [72]. For example, blocking p75 by short-hairpin RNA in cultured Lewis lung carcinoma cells results in decreased TNF-mediated expression of VEGF, placental growth factor and HGF, suggesting that p75 is essential factor for tumor angiogenesis [72]. Similarly, blocking TNF- α inhibits angiogenesis in metastatic oral squamous cell carcinoma cells (sh-IFIT2 meta cell) in NOD/SCID mice [90]. TGF- β negatively regulates VEGF-A expression via a PKA- and Smad2-independent and Smad3-dependent pathways as demonstrated in FET α /DNRII colon cancer cell lines [73]. HGF is an angiogenic factor secreted predominantly by fibroblasts; it stimulates invasiveness of cancer cells via c-Met receptor tyrosine kinase activation [79, 91, 92]. In fact, high HGF serum levels is correlated with VEGF and IL-8 expression, advanced tumor stage and poor survival of esophageal squamous cell carcinoma (ESCC) patients [78]. High expression of another pro-angiogenic factor, EGFR correlates with increased microvessel density resulting in enhanced tumor angiogenesis via the HIF-1 α and Notch1 pathways in tissues from head and neck squamous cell carcinoma patients [76]. Neuropilin is a transmembrane glycoprotein which serves as a receptors or co-receptor for multiple ligands including VEGF, HGF, EGF and FGF which are involved in tumor angiogenesis [93, 94]. In gastric cancer, high expression of neuropilin correlates with advanced clinical stages (III and IV) [95]. Depletion of neuropilin-1 inhibits the activation of EGF/EGFR, VEGF/VEGFR2 and HGF/c-Met angiogenic pathways activated by recombinant human VEGF-165, HGF and EGF proteins [91, 95]. Thus, the role of neurotrophic factors such as NGF, BDNF and their molecular pathways should be considered in the development of anti-angiogenic therapies.

Concluding Remarks

Despite the increasing interest to the role of the nervous system in cancer development and progression, the knowledge in this area is scarce. Most neurotransmitters released by nerve fibers promote tumor angiogenesis, however, some neurotransmitters induce anti-cancer effects. Whether these effects are cancer type or receptor dependent need further elucidation.

To date, most studies investigating the role of the nervous system in modulation of tumor angiogenesis have been performed in cell lines and animal models. Limited studies are available from cancer patients and at different stages of disease. Understanding molecular mechanisms by which nervous system modulates tumor angiogenesis may open new avenues for understanding mechanisms of tumor angiogenesis, identification of new biomarkers for cancer diagnosis and prognosis, and, defining novel targets for therapeutic interventions.

Acknowledgements NK was supported by an Australian Postgraduate Research Award, LS and KN was supported by the College of Health and Biomedicine Victoria University, Australia and VA was supported by the Centre for Chronic Disease, Victoria University, Australia.

Compliance with Ethical Standards

Conflict of Interest The authors confirm that this article content has not conflict of interest.

References

1. Folkman J (2007) Angiogenesis: an organizing principle for drug discovery? *Nat Rev Drug Discov* 6(4):273–286
2. Zhao Y, Adjei AA (2015) Targeting angiogenesis in cancer therapy: moving beyond vascular endothelial growth factor. *Oncologist* 20(6):660–673
3. Nishida N, Yano H, Nishida T, Kamura T, Kojiro M (2006) Angiogenesis in cancer. *Vasc Health Risk Manag* 2(3):213–219
4. McMahon G (2000) VEGF receptor signaling in tumor angiogenesis. *Oncologist* 5(Supplement 1):3–10
5. Goel S, Duda DG, Xu L, Munn LL, Boucher Y, Fukumura D, Jain RK (2011) Normalization of the vasculature for treatment of cancer and other diseases. *Physiol Rev* 91(3):1071–1121
6. Zibara K, Awada Z, Dib L, El-Saghir J, Al-Ghadban S, Ibrik A, El-Zein N, El-Sabban M (2015) Anti-angiogenesis therapy and gap junction inhibition reduce MDA-MB-231 breast cancer cell invasion and metastasis in vitro and in vivo. *Sci Rep* 5:12598
7. Rusckowski M, Wang Y, Blankenberg FG, Levashova Z, Backer MV, Backer JM (2016) Targeted scVEGF/177Lu radiopharmaceutical inhibits growth of metastases and can be effectively combined with chemotherapy. *EJNMMI Res* 6(1):1–9
8. Vasudev NS, Reynolds AR (2014) Anti-angiogenic therapy for cancer: current progress, unresolved questions and future directions. *Angiogenesis* 17(3):471–494
9. Quail DF, Joyce JA (2013) Microenvironmental regulation of tumor progression and metastasis. *Nat Med* 19(11):1423–1437
10. Gupta GP, Massagué J (2006) Cancer Metastasis: building a framework. *Cell* 127(4):679–695

11. Place AE, Jin Huh S, Polyak K (2011) The microenvironment in breast cancer progression: biology and implications for treatment. *Breast Cancer Res* 13(6):227
12. Weis SM, Cheresch DA (2011) Tumor angiogenesis: molecular pathways and therapeutic targets. *Nat Med* 17(11):1359–1370
13. Jobling P, Pundavela J, Oliveira SM, Roselli S, Walker MM, Hondemarck H (2015) Nerve–cancer cell cross-talk: a novel promoter of tumor progression. *Cancer Res* 75(9):1777–1781
14. Ondicova K, Mravec B (2010) Role of nervous system in cancer aetiopathogenesis. *Lancet Oncol* 11(6):596–601
15. Chen H, Liu D, Yang Z, Sun L, Deng Q, Yang S, Qian L, Guo L, Yu M, Hu M, Shi M, Guo N (2014) Adrenergic signaling promotes angiogenesis through endothelial cell-tumor cell crosstalk. *Endocr Relat Cancer* 21(5):783–795
16. Liu J, Deng G-H, Zhang J, Wang Y, Xia X-Y, Luo X-M, Deng Y-T, He S-S, Mao Y-Y, Peng X-C (2015) The effect of chronic stress on anti-angiogenesis of sunitinib in colorectal cancer models. *Psychoneuroendocrinology* 52:130–142
17. Deng G-H, Liu J, Zhang J, Wang Y, Peng X-C, Wei Y-Q, Jiang Y (2014) Exogenous norepinephrine attenuates the efficacy of sunitinib in a mouse cancer model. *J Exp Clin Cancer Res* 33(1):1–12
18. Moreno-Smith M, Lee SJ, Lu C, Nagaraja AS, He G, Rupaimoole R, Han HD, Jennings NB, Roh J-W, Nishimura M (2013) Biologic effects of dopamine on tumor vasculature in ovarian carcinoma. *Neoplasia* 15(5):502–IN515
19. Moreno-Smith M, Lu C, Shahzad MM, Pena GNA, Allen JK, Stone RL, Mangala LS, Han HD, Kim HS, Farley D (2011) Dopamine blocks stress-mediated ovarian carcinoma growth. *Clin Cancer Res* 17(11):3649–3659
20. Chakroborty D, Sarkar C, Mitra RB, Banerjee S, Dasgupta PS, Basu S (2004) Depleted dopamine in gastric cancer tissues: dopamine treatment retards growth of gastric cancer by inhibiting angiogenesis. *Clin Cancer Res* 10(13):4349–4356
21. Hoepfner LH, Wang Y, Sharma A, Javeed N, Van Keulen VP, Wang E, Yang P, Roden AC, Peikert T, Molina JR (2015) Dopamine D2 receptor agonists inhibit lung cancer progression by reducing angiogenesis and tumor infiltrating myeloid derived suppressor cells. *Mol Oncol* 9(1):270–281
22. Fava G, Marucci L, Glaser S, Francis H, De Morrow S, Benedetti A, Alvaro D, Venter J, Meininger C, Patel T (2005) γ -aminobutyric acid inhibits cholangiocarcinoma growth by cyclic AMP-dependent regulation of the protein kinase a/extracellular signal-regulated kinase 1/2 pathway. *Cancer Res* 65(24):11437–11446
23. Park SH, Kim B-R, Lee JH, Park ST, Lee S-H, Dong SM, Rho SB (2014) GABARBP down-regulates HIF-1 α expression through the VEGFR-2 and PI3K/mTOR/4E-BP1 pathways. *Cell Signal* 26(7):1506–1513
24. Nocito A, Dahm F, Jochum W, Jang JH, Georgiev P, Bader M, Graf R, Clavien P-A (2008) Serotonin regulates macrophage-mediated angiogenesis in a mouse model of colon cancer allografts. *Cancer Res* 68(13):5152–5158
25. Speyer CL, Hachem AH, Assi A, DeVries JA, Gorski DH (2013) Metabotropic glutamate receptor-1 as a novel target for the anti-angiogenic treatment of breast cancer. *Cancer Res* 73(8 Supplement):3895
26. Wen Y, Li J, Koo J, Shin SS, Lin Y, Jeong BS, Mehnert JM, Chen S, Cohen-Sola KA, Goydos JS (2014) Activation of the glutamate receptor GRM1 enhances angiogenic signaling to drive melanoma progression. *Cancer Res* 74(9):2499–2509
27. Zhang Q, Tang X, Zhang Z-F, Velikina R, Shi S, Le AD (2007) Nicotine induces hypoxia-inducible factor-1 α expression in human lung cancer cells via nicotinic acetylcholine receptor-mediated signaling pathways. *Clin Cancer Res* 13(16):4686–4694
28. Wong HPS, Yu L, Lam EKY, Tai EKK, Wu WKK, Cho C-H (2007) Nicotine promotes colon tumor growth and angiogenesis through β -adrenergic activation. *Toxicol Sci* 97(2):279–287
29. Lombardi Mí G, Negroni M, Pelegrina LT, Castro Mí E, Fiszman GL, Azar Mí E, Morgado CC, Sales Mí E (2013) Autoantibodies against muscarinic receptors in breast cancer: their role in tumor angiogenesis. *PLoS One* 8(2):e57572
30. de la Torre E, Davel L, Jansis MA, Gotoh T, de Lustig ES, Sales ME (2005) Muscarinic receptors participation in angiogenic response induced by macrophages from mammary adenocarcinoma-bearing mice. *Breast Cancer Res* 7(3):R345–R352
31. Medeiros PJ, Jackson DN (2013) Neuropeptide Y Y5-receptor activation on breast cancer cells acts as a paracrine system that stimulates VEGF expression and secretion to promote angiogenesis. *Peptides* 48:106–113
32. Medeiros PJ, Al-Khazraji BK, Novielli NM, Postovit LM, Chambers AF, Jackson DN, Neuropeptide Y (2012) Stimulates proliferation and migration in the 4T1 breast cancer cell line. *Int J Cancer J Int du Cancer* 131(2):276–286
33. Alasvand M, Rashidi B, Javanmard SH, Akhavan MM, Khazaei M (2015) Effect of blocking of neuropeptide Y Y2 receptor on tumor angiogenesis and progression in normal and diet-induced obese C57BL/6 mice. *Glob J Health Sci* 7(7):46883
34. Lu C, Everhart L, Tilan J, Kuo L, Sun CJ, Munivenkatappa RB, Jönsson-Rylander A-C, Sun J, Kuan-Celariar A, Li L (2010) Neuropeptide Y and its Y2 receptor: potential targets in neuroblastoma therapy. *Oncogene* 29(41):5630–5642
35. Nakamura Y, Yasuoka H, Tsujimoto M, Yoshidome K, Nakahara M, Nakao K, Nakamura M, Kakudo K (2006) Nitric oxide in breast cancer: induction of vascular endothelial growth factor-C and correlation with metastasis and poor prognosis. *Clin Cancer Res* 12(4):1201–1207
36. Nomellini RS, Ribeiro LCDA, Tavares-Murta BM, Adad SJ, Murta EFC (2009) Production of nitric oxide and expression of inducible nitric oxide synthase in ovarian cystic tumors. *Mediators Inflamm* 2008:186584
37. Wang L, Shi GG, Yao JC, Gong W, Wei D, Wu TT, Ajani JA, Huang S, Xie K (2005) Expression of endothelial nitric oxide synthase correlates with the angiogenic phenotype of and predicts poor prognosis in human gastric cancer. *Gastric Cancer* 8(1):18–28
38. Sarkar C, Chakroborty D, Basu S (2013) Neurotransmitters as regulators of tumor angiogenesis and immunity: the role of catecholamines. *J Neuroimmune Pharmacol Off J Soc NeuroImmune Pharmacol* 8(1):7–14
39. Chakroborty D, Sarkar C, Basu B, Dasgupta PS, Basu S (2009) Catecholamines regulate tumor angiogenesis. *Cancer Res* 69(9):3727–3730
40. Thaker PH, Han LY, Kamat AA, Arevalo JM, Takahashi R, Lu C, Jennings NB, Armaiz-Pena G, Bankson JA, Ravoory M (2006) Chronic stress promotes tumor growth and angiogenesis in a mouse model of ovarian carcinoma. *Nat Med* 12(8):939–944
41. Xie H, Li C, He Y, Griffin R, Ye Q, Li L (2015) Chronic stress promotes oral cancer growth and angiogenesis with increased circulating catecholamine and glucocorticoid levels in a mouse model. *Oral Oncol* 51(11):991–997
42. Sloan EK, Priceman SJ, Cox BF, Yu S, Pimentel MA, Tanganangnukul V, Arevalo JM, Morizono K, Karanikolas BD, Wu L (2010) The sympathetic nervous system induces a metastatic switch in primary breast cancer. *Cancer Res* 70(18):7042–7052
43. Madden KS, Szpunar MJ, Brown EB (2011) Beta-adrenergic receptors (beta-AR) regulate VEGF and IL-6 production by divergent pathways in high beta-AR-expressing breast cancer cell lines. *Breast Cancer Res Treat* 130(3):747–758
44. Reedijk M, Odorcic S, Chang L, Zhang H, Miller N, McCready DR, Lockwood G, Egan SE (2005) High-level coexpression of Jag1 and Notch1 is observed in human breast cancer and is associated with poor overall survival. *Cancer Res* 65(18):8530–8537

45. Dickson BC, Mulligan AM, Zhang H, Lockwood G, O'Malley FP, Egan SE, Reedijk M (2007) High-level JAG1 mRNA and protein predict poor outcome in breast cancer. *Mod Pathol* 20(6):685–693
46. Basu S, Nagy JA, Pal S, Vasile E, Eckelhoefer IA, Bliss VS, Manseau EJ, Dasgupta PS, Dvorak HF, Mukhopadhyay D (2001) The neurotransmitter dopamine inhibits angiogenesis induced by vascular permeability factor/vascular endothelial growth factor. *Nat Med* 7(5):569–574
47. Tilan J, Kitlinska J (2010) Sympathetic neurotransmitters and tumor angiogenesis-link between stress and cancer progression. *J Oncol* 2010:539706
48. Sarkar C, Chakroborty D, Chowdhury UR, Dasgupta PS, Basu S (2008) Dopamine increases the efficacy of anticancer drugs in breast and colon cancer preclinical models. *Clin Cancer Res* 14(8):2502–2510
49. Schuller HM (2009) Is cancer triggered by altered signalling of nicotinic acetylcholine receptors? *Nat Rev Cancer* 9(3):195–205
50. Egleton RD, Brown KC, Dasgupta P (2009) Angiogenic activity of nicotinic acetylcholine receptors: implications in tobacco-related vascular diseases. *Pharmacol Ther* 121(2):205–223
51. Cucina A, Sapienza P, Corvino V, Borrelli V, Mariani V, Randone B, Santoro D'Angelo L, Cavallaro A (2000) Nicotine-induced smooth muscle cell proliferation is mediated through bFGF and TGF-beta 1. *Surgery* 127(3):316–322
52. Conklin BS, Zhao W, Zhong DS, Chen C (2002) Nicotine and cotinine up-regulate vascular endothelial growth factor expression in endothelial cells. *Am J Pathol* 160(2):413–418
53. Brown KC, Lau JK, Dom AM, Witte TR, Luo H, Crabtree CM, Shah YH, Shiflett BS, Marcelo AJ, Proper NA, Hardman WE, Egleton RD, Chen YC, Mangiarua EI, Dasgupta P (2012) MG624, an alpha7-nAChR antagonist, inhibits angiogenesis via the Egr-1/FGF2 pathway. *Angiogenesis* 15(1):99–114
54. Davis R, Rizwani W, Banerjee S, Kovacs M, Haura E, Coppola D, Chellappan S (2009) Nicotine promotes tumor growth and metastasis in mouse models of lung cancer. *PLoS One* 4(10):e7524
55. Lee CH, Huang CS, Chen CS, Tu SH, Wang YJ, Chang YJ, Tam KW, Wei PL, Cheng TC, Chu JS, Chen LC, Wu CH, Ho YS (2010) Overexpression and activation of the alpha9-nicotinic receptor during tumorigenesis in human breast epithelial cells. *J Natl Cancer Inst* 102(17):1322–1335
56. Wu C-H, Lee C-H, Ho Y-S (2011) Nicotinic acetylcholine receptor-based blockade: applications of molecular targets for cancer therapy. *Clin Cancer Res* 17(11):3533–3541
57. Dasgupta P, Chellappan SP (2006) Nicotine-mediated cell proliferation and angiogenesis: new twists to an old story. *Cell Cycle* 5(20):2324–2328
58. Fiszman GL, Middonno MC, de la Torre E, Farina M, Español AJ, Sales ME (2007) Activation of muscarinic cholinergic receptors induces MCF-7 cells proliferation and angiogenesis by stimulating nitric oxide synthase activity. *Cancer Biol Ther* 6(7):1106–1113
59. Li L, Kilbourn RG, Adams J, Fidler IJ (1991) Role of nitric oxide in lysis of tumor cells by cytokine-activated endothelial cells. *Cancer Res* 51(10):2531–2535
60. Cianchi F, Cortesini C, Fantappiè O, Messerini L, Schiavone N, Vannacci A, Nistri S, Sardi I, Baroni G, Marzocca C, Perna F, Mazzanti R, Becchi P, Masini E (2003) Inducible nitric oxide synthase expression in human colorectal cancer: correlation with tumor angiogenesis. *Am J Pathol* 162(3):793–801
61. Fukumura D, Kashiwagi S, Jain RK (2006) The role of nitric oxide in tumour progression. *Nat Rev Cancer* 6(7):521–534
62. Ridnour LA, Thomas DD, Donzelli S, Espey MG, Roberts DD, Wink DA, Isenberg JS (2006) The biphasic nature of nitric oxide responses in tumor biology. *Antioxid Redox Signal* 8(7–8):1329–1337
63. Andrade SP, Hart IR, Piper PJ (1992) Inhibitors of nitric oxide synthase selectively reduce flow in tumour-associated neovasculature. *Br J Pharmacol* 107(4):1092–1095
64. Chu SC, Marks-Konczalik J, Wu H-P, Banks TC, Moss J (1998) Analysis of the cytokine-stimulated human inducible nitric oxide synthase (iNOS) gene: characterization of differences between human and mouse iNOS promoters. *Biochem Biophys Res Commun* 248(3):871–878
65. Vahora H, Khan MA, Alalami U, Hussain A (2016) The potential role of nitric oxide in halting cancer progression through chemoprevention. *J Cancer Prev* 21(1):1–12
66. Radisky ES, Radisky DC (2015) Matrix metalloproteinases as breast cancer drivers and therapeutic targets. *Front Biosci (Landmark ed)* 20:1144–1163
67. Wangari-Talbot J, Wall BA, Goydos JS, Chen S (2012) Functional effects of GRM1 suppression in human melanoma cells. *Mol Cancer Res* 10(11):1440–1450
68. Xia H, Zhao YN, Yu CH, Zhao YL, Liu Y (2016) Inhibition of metabotropic glutamate receptor 1 suppresses tumor growth and angiogenesis in experimental non-small cell lung cancer. *Eur J Pharmacol* 783:103–111
69. He T, Qi F, Jia L, Wang S, Wang C, Song N, Fu Y, Li L, Luo Y (2015) Tumor cell-secreted angiogenin induces angiogenic activity of endothelial cells by suppressing miR-542-3p. *Cancer Lett* 368(1):115–125
70. Miyake M, Goodison S, Lawton A, Gomes-Giacoa E, Rosser C (2015) Angiogenin promotes tumoral growth and angiogenesis by regulating matrix metalloproteinase-2 expression via the ERK1/2 pathway. *Oncogene* 34(7):890–901
71. Shu J, Huang M, Tian Q, Shui Q, Zhou Y, Chen J (2015) Downregulation of angiogenin inhibits the growth and induces apoptosis in human bladder cancer cells through regulating AKT/mTOR signaling pathway. *J Mol Histol* 46(2):157–171
72. Sasi SP, Yan X, Enderling H, Park D, Gilbert H-Y, Curry C, Coleman C, Hlatky L, Qin G, Kishore R (2012) Breaking the 'harmony' of TNF- α signaling for cancer treatment. *Oncogene* 31(37):4117–4127
73. Geng L, Chaudhuri A, Talmon G, Wisecarver JL, Wang J (2013) TGF-Beta suppresses VEGFA-mediated angiogenesis in colon cancer metastasis. *PLoS One* 8(3):e59918
74. Lin C-Y, Hung S-Y, Chen H-T, Tsou H-K, Fong Y-C, Wang S-W, Tang C-H (2014) Brain-derived neurotrophic factor increases vascular endothelial growth factor expression and enhances angiogenesis in human chondrosarcoma cells. *Biochem Pharmacol* 91(4):522–533
75. Larrieu-Lahargue F, Welm AL, Bouchecareilh M, Alitalo K, Li DY, Bikfalvi A, Auguste P (2012) Blocking fibroblast growth factor receptor signaling inhibits tumor growth, lymphangiogenesis, and metastasis. *PLoS One* 7(6):e39540
76. Wang W-M, Zhao Z-L, Ma S-R, Yu G-T, Liu B, Zhang L, Zhang W-F, Kulkarni AB, Sun Z-J, Zhao Y-F (2015) Epidermal growth factor receptor inhibition reduces angiogenesis via hypoxia-inducible factor-1 α and notch1 in head neck squamous cell carcinoma. *PLoS One* 10(2):e0119723
77. Romon R, Adriaenssens E, Lagadec C, Germain E, Hondermarck H, Le Bourhis X (2010) Nerve growth factor promotes breast cancer angiogenesis by activating multiple pathways. *Breast Cancer* 9:11
78. Ren Y, Cao B, Law S, Xie Y, Lee PY, Cheung L, Chen Y, Huang X, Chan HM, Zhao P (2005) Hepatocyte growth factor promotes cancer cell migration and angiogenic factors expression: a prognostic marker of human esophageal squamous cell carcinomas. *Clin Cancer Res* 11(17):6190–6197
79. Sugie S, Mukai S, Yamasaki K, Kamibeppu T, Tsukino H, Kamoto T (2016) Plasma macrophage-stimulating protein and hepatocyte

- growth factor levels are associated with prostate cancer progression. *Hum Cell* 29(1):22–29
80. Walsh EM, Kim R, Del Valle L, Weaver M, Sheffield J, Lazarovici P, Marcinkiewicz C (2012) Importance of interaction between nerve growth factor and $\alpha 9 \beta 1$ integrin in glial tumor angiogenesis. *Neuro-Oncology* 14(7):890–901
 81. Staniszewska I, Zaveri S, Del Valle L, Oliva I, Rothman VL, Croul SE, Roberts DD, Mosher DF, Tuszynski GP, Marcinkiewicz C (2007) Interaction of $\alpha 9 \beta 1$ integrin with thrombospondin-1 promotes angiogenesis. *Circ Res* 100(9):1308–1316
 82. Vlahakis NE, Young BA, Atakilit A, Hawkrigde AE, Issaka RB, Boudreau N, Sheppard D (2007) Integrin $\alpha 9 \beta 1$ directly binds to vascular endothelial growth factor (VEGF)-a and contributes to VEGF-A-induced angiogenesis. *J Biol Chem* 282(20):15187–15196
 83. Staniszewska I, Sariyer IK, Lecht S, Brown MC, Walsh EM, Tuszynski GP, Safak M, Lazarovici P, Marcinkiewicz C (2008) Integrin $\alpha 9 \beta 1$ is a receptor for nerve growth factor and other neurotrophins. *J Cell Sci* 121(4):504–513
 84. Nilsson UW, Abrahamsson A, Dabrosin C (2010) Angiogenin regulation by estradiol in breast tissue: tamoxifen inhibits angiogenin nuclear translocation and antiangiogenin therapy reduces breast cancer growth in vivo. *Clin Cancer Res* 16(14):3659–3669
 85. Katona TM, Neubauer BL, Iversen PW, Zhang S, Baldrige LA, Cheng L (2005) Elevated expression of angiogenin in prostate cancer and its precursors. *Clin Cancer Res* 11(23):8358–8363
 86. Hisai H, Kato J, Kobune M, Murakami T, Miyanishi K, Takahashi M, Yoshizaki N, Takimoto R, Terui T, Niitsu Y (2003) Increased expression of angiogenin in hepatocellular carcinoma in correlation with tumor vascularity. *Clin Cancer Res* 9(13):4852–4859
 87. Miyake H, Hara I, Yamanaka K, Gohji K, Arakawa S, Kamidono S (1999) Increased angiogenin expression in the tumor tissue and serum of urothelial carcinoma patients is related to disease progression and recurrence. *Cancer* 86(2):316–324
 88. Gao X, Xu Z (2008) Mechanisms of action of angiogenin. *Acta Biochim Biophys Sin* 40(7):619–624
 89. de Aguiar RB, Parise CB, Souza CRT, Braggion C, Quintilio W, Moro AM, Navarro Marques FL, Buchpiguel CA, Chammas R, de Moraes JZ (2016) Blocking FGF2 with a new specific monoclonal antibody impairs angiogenesis and experimental metastatic melanoma, suggesting a potential role in adjuvant settings. *Cancer Lett* 371(2):151–160
 90. Lai KC, Liu CJ, Lin TJ, Mar AC, Wang HH, Chen CW, Hong ZX, Lee TC (2016) Blocking TNF-alpha inhibits angiogenesis and growth of IFIT2-depleted metastatic oral squamous cell carcinoma cells. *Cancer Lett* 370(2):207–215
 91. Kawaguchi M, Kataoka H (2014) Mechanisms of hepatocyte growth factor activation in cancer tissues. *Cancers* 6(4):1890–1904
 92. Kataoka H, Miyata S, Uchinokura S, Itoh H (2003) Roles of hepatocyte growth factor (HGF) activator and HGF activator inhibitor in the pericellular activation of HGF/scatter factor. *Cancer Metastasis Rev* 22(2–3):223–236
 93. Ellis LM (2006) The role of neuropilins in cancer. *Mol Cancer Ther* 5(5):1099–1107
 94. Djordjevic S, Driscoll PC (2013) Targeting VEGF signalling via the neuropilin co-receptor. *Drug Discov Today* 18(9–10):447–455
 95. Li L, Jiang X, Zhang Q, Dong X, Gao Y, He Y, Qiao H, Xie F, Xie X, Sun X (2016) Neuropilin-1 is associated with clinicopathology of gastric cancer and contributes to cell proliferation and migration as multifunctional co-receptors. *J Exp Clin Cancer Res* 35(1):16